

AMERICAN SOCIETY FOR REPRODUCTIVE MEDICINE

MULTIPLE PREGNANCY
AND BIRTH:
TWINS, TRIPLETS,
& HIGHER ORDER MULTIPLES

A Guide for Patients



PATIENT INFORMATION SERIES

Published by the American Society for Reproductive Medicine under the direction of the Patient Education Committee and the Publications Committee. No portion herein may be reproduced in any form without written permission. This booklet is in no way intended to replace, dictate, or fully define evaluation and treatment by a qualified physician. It is intended solely as an aid for patients seeking general information on issues in reproductive medicine.

Copyright 2004 by American Society for Reproductive Medicine.

MULTIPLE PREGNANCY AND BIRTH: TWINS, TRIPLETS, & HIGHER ORDER MULTIPLES

A Guide for Patients

A glossary of italicized words is located at the end of this booklet.

INTRODUCTION

Multiple births are much more common today than they were in the past. According to the U.S. Department of Health and Human Services, the twin birth rate has increased over 50% since 1980, and triplet, quadruplet, and higher order multiple births have increased at an even higher rate. There are more multiple births today in part because more women are receiving *infertility* treatment, which carries a risk of multiple pregnancy. Also, more women are waiting until later in life to attempt pregnancy, and older women are more likely than younger women to get pregnant with multiples, especially with fertility treatment. Although major medical advances have improved the outcomes of multiple births, they are still associated with significant medical risks and complications for the mother and children. If you are at risk for a multiple pregnancy, this booklet will help you learn how and why multiple pregnancies occur, and the unique issues associated with carrying and delivering a multiple pregnancy.

TWINS- THE MOST COMMON MULTIPLE

You may know someone who has twins, but do you know how twins occur and how they develop?


There are two types of twins: identical and fraternal (non-identical).

Identical twins occur when a single *embryo*, created by the union of a *sperm*

and egg, divides into two embryos. Each embryo is *monozygotic*, genetically identical. When the twinning process occurs early in the pregnancy, in the first two weeks after conception, each *fetus* has its own separate *gestational sac* and *placenta*. However, if the twinning process occurs later, after the placenta has formed, the two embryos may be together in one sac.

Non-identical twins occur when two separate eggs are each fertilized by a separate sperm. The two embryos that result are *dizygotic*, not genetically identical. Therefore, the twins are not identical.

If *ultrasound* shows twins together in a single sac, they are identical. Twins that are in the same sac but separated by a thin membrane, the *amnion*, are also identical. Twins in two different sacs may be identical or non-identical.



The “Vanishing Twin Syndrome”

Sometimes, very early in a twin pregnancy, one of the fetuses “disappears.” This is referred to as “vanishing twin syndrome.” Even after ultrasound has shown heart movement in twins, spontaneous loss of one of the fetuses occurs in up to 20% of twin pregnancies. Spontaneous losses are even higher in triplet and quadruplet pregnancies. A fetal loss rate of 40% may occur in pregnancies with triplets or more. When a fetus is lost in the first trimester, the remaining fetus or fetuses generally continue to develop normally, although vaginal bleeding may occur.

Ultrasound examinations performed early in the fifth week of pregnancy may occasionally fail to identify all fetuses. An “appearing twin” may be found after the fifth week in nearly 10% of non-identical twin or multiple pregnancies and over 80% of cases of identical twins. After six to eight weeks, ultrasound should provide an accurate assessment for the number of fetuses.

Risk Factors for Multiple Pregnancy

Naturally, twins occur in about one in 250 pregnancies, triplets in about one in 8,000 pregnancies, and quadruplets in about one in 700,000 pregnancies. The main factor that increases your chances of having a multiple pregnancy is the use of infertility treatment, but there are other factors

Your race, age, heredity, or history of prior pregnancy does not increase your chances of having identical twins, but does increase your chance of having non-identical twins. Infertility treatment increases your risk of having twins, both identical and non-identical.

Race. Twins occur in approximately 1 of every 90 pregnancies in North America. The incidence is higher in Africa, with a rate of 1 in 20 births in Nigeria. Twins are less common in Asia. In Japan, for example, twins occur only once in every 155 births.

Heredity. The mother's family history may be more significant than the father's. Non-identical twin women give birth to twins at the rate of 1 set per 60 births. However, non-identical males father twins at a rate of 1 set per 125 births.

Maternal age and prior pregnancy history. The frequency of twins increases with maternal age and number of pregnancies. Women between 35 to 40 years of age with four or more children are three times more likely to have twins than a woman under 20 without children.

Maternal height and weight. Non-identical twins are more common in large and tall women than in small women. This may be related more to nutrition than to body size alone. During World War II, the incidence of non-identical twinning decreased in Europe when food was not readily available.

Fertility Drugs and Assisted Reproductive Technology

Multiple pregnancy is more common in women who utilize fertility medications to undergo *ovulation induction* or *superovulation*. Of women who achieve pregnancy with *clomiphene citrate*, approximately 5% to 12% are twins, and less than 1% are triplets or greater.

Approximately 20% of pregnancies resulting from *gonadotropins* are multiples. While most of these pregnancies are twins, up to 5% are triplets or greater due to the release of more eggs than expected.

In part, ART procedures such as *in vitro fertilization* (IVF) are largely responsible for the increase in the multiple birth rate. The risk of multiple pregnancy increases as the number of embryos transferred increases. Superovulation accounts for a comparable proportion of high order multiples.

DURATION OF MULTIPLE PREGNANCIES

The duration of a normal *singleton* pregnancy ranges from 37 weeks to 42 weeks from the time of the last menstrual period. Twin pregnancies occasionally progress to 40 weeks but almost always deliver early. As the number of fetuses increases, the expected duration of the pregnancy decreases. The average duration is 35 weeks for twins, 33 weeks for triplets, and 29 weeks for quadruplets.

| TYPE OF PREGNANCY | AVERAGE GESTATIONAL AGE AT TIME OF DELIVERY | AVERAGE BIRTH WEIGHT |
|-------------------|---|------------------------|
| Singleton | 40 weeks | 7 lbs. (3,300 grams) |
| Twin | 35 weeks | 5.5 lbs. (2,500 grams) |
| Triplet | 33 weeks | 4 lbs. (1,800 grams) |
| Quadruplets | 29 weeks | 3 lbs. (1,400 grams) |

COMPLICATIONS OF MULTIPLE PREGNANCIES

Complications increase with each additional fetus in a multiple pregnancy and include severe nausea and vomiting, Cesarean section, or forceps delivery. If you are pregnant with twins or more, or if you are at risk for a multiple pregnancy, you should be aware of these and other potential problems you might experience.

Premature Birth

Premature labor and delivery pose the greatest risk to a multiple pregnancy. Feasibility of a vaginal delivery depends on the size, position, and health of the infants, as well as the size and shape of the pelvic bones. Cesarean section is often needed for twin pregnancies and is expected for delivery of triplets.

Although only 1% of all deliveries are twins, they account for 10% of all premature deliveries. Compared to singletons, of which eight per 1,000 die in the first month of life, twins are seven times more likely to die, and triplets are 20 times more likely to die.

Since premature labor and delivery present such a serious risk, the pregnant mother must understand the warning signs for early labor. Pelvic pressure, low back pain, increased vaginal discharge, or a change in the frequency of "false-labor" pains should be reported to the physician, who can sometimes delay premature delivery by a few days or more if it is detected early. Each day gained provides valuable fetal growth and development.

Once advanced labor is established, delivery cannot be stopped. In rare instances, delivery of a second twin can be delayed. This delay, when possible, allows for continued growth in the protective environment of the *uterus*.

Placental Problems

The placenta is attached to the wall of the uterus, and the fetus is attached to the placenta by the umbilical cord. The placenta provides blood, oxygen, and nutrition to the fetus through the umbilical cord.

Placental function is likely to be abnormal in a multiple pregnancy. The placenta ages prematurely and may slow fetal growth, especially late in the third trimester. If the placenta is unable to provide adequate oxygen or nutrients to the fetus, the fetus cannot grow properly. Twins and multiples that are more than 30% “underweight” by ultrasound measurements are at increased risk of complications and have death rates of nearly 25%.

Another placental problem is twin-twin transfusion, a life threatening condition in identical twins. This transfusion occurs when blood flows from one fetus to the other. Poor growth occurs in the “donor” twin, and excessive blood passes to the “recipient” twin. Therapeutic *amniocentesis* and laser coagulation of blood vessels may reduce complications of twin-twin transfusion.

Preeclampsia

Preeclampsia, also known as *toxemia*, occurs three to five times more often in multiple pregnancies. *Preeclampsia* is diagnosed when the mother’s blood pressure becomes elevated and protein is detected in the urine. The condition may progress and threaten the health of the mother and the pregnancy. When severe, the mother may have seizures or even a stroke.

Diabetes

Women with multiple pregnancies are more likely to develop *gestational diabetes* during pregnancy. Babies of diabetic mothers are more likely to experience respiratory distress and other newborn complications.

Fetal and Newborn Complications

Premature delivery places an infant at increased risk for severe complications or early death. A baby’s lungs, brain, circulatory system, intestinal system, and eyes may be too immature. Survivors of premature birth may have lifelong handicaps.

Of the premature babies who die, 50% succumb to *respiratory distress syndrome*, the inability to circulate oxygen from the lungs throughout the body. Brain damage is responsible for almost 10% of premature newborn deaths. Birth defects and stillbirths account for about 30% of the deaths in twins and multiple pregnancies.

Low birth weight of less than 5 and a half pounds (2,500 grams) occurs in 50% of twins. The average birth weight is approximately 4 pounds (1,800

grams) for triplets and 3 pounds (1,400 grams) for quadruplets. As a result of prematurity, the risk for *cerebral palsy* is four times more likely to occur in twins. The rate is even greater for triplets and higher order multiple births. Prematurity may also result in visual impairment or blindness.

The overall survival rate is 85% for newborns over 2 lb., 3 oz. (1,000 grams), and less than 40% for those under 2 lb., 3 oz. Birth weight also corresponds closely to the severity of disability throughout the childhood years. Disability occurs in over 25% of children with a birth weight less than 2 lb., 3 oz.

PREVENTION OF MULTIPLE PREGNANCY

Prevention during infertility treatment is the best approach to avoiding a multiple pregnancy. In ART cycles, limiting the number of embryos transferred is an effective approach. Consult the ASRM Practice Committee Report titled “Guidelines on Number of Embryos Transferred” for recommendations regarding the optimal number of embryos to transfer based on patient age, embryo quality, and other criteria.

In the United States, physicians and patients jointly decide how many embryos to transfer. However, in England, no more than two embryos may be transferred in most cases. In Canada, a maximum of three embryos are recommended for transfer.

The ultimate goal of physicians performing ART is to achieve a high pregnancy rate while transferring a single embryo. While physicians can transfer two embryos and still maintain acceptable pregnancy rates, the transfer of one embryo may eventually be associated with good pregnancy rates, thereby resolving the problem of multiple pregnancies caused by multiple embryo transfer.

Multiple pregnancies are a known complication of ovulation drugs. Most physicians monitor patients with ultrasound examinations and blood tests. A woman with a large number of ovarian *follicles* or high *hormone* levels has an increased risk of a multiple pregnancy, and the ART cycle may be canceled to avoid the risk.

Multifetal Pregnancy Reduction

When a triplet or higher order multiple pregnancy occurs, *multifetal pregnancy reduction* may be considered for the health of the mother and to improve survival of the pregnancy. While multifetal pregnancy reduction carries some risk of a complete *miscarriage*, it also reduces the chances of extreme premature birth. For more information, see the ASRM Patient Fact Sheets *Multiple Gestation & Multifetal Pregnancy Reduction* and *Challenges of Parenting Multiples*.

CARRYING A MULTIPLE PREGNANCY

In order to achieve the best outcome with a multiple pregnancy, the expectant mother must work as part of the health care team. A nearly total change in lifestyle can be expected, especially after about 20 weeks into the pregnancy.

Metabolic and Nutritional Considerations

There is an increased need for maternal nutrition in multiple pregnancies. An expectant mother needs to gain more weight in a multiple pregnancy, especially if she begins the pregnancy underweight. With multiples, weight gain of approximately 45 pounds is optimal for normal weight women. The pattern of weight gain is important too. Healthy birth weights are most likely achieved when the mother gains nearly one pound per week in the first 20 weeks. In addition to weight gain, most physicians recommend increasing vitamin supplementation by 50% to 100%. The increase in fetal growth with appropriate nutrition and weight gain may greatly improve pregnancy outcome at a minimum of cost.

Activity Precautions

Many physicians who manage multiple pregnancies believe that a reduction in activities and increased bed rest prolongs these pregnancies and improves outcomes. Women with multiple pregnancies are usually advised to avoid strenuous activity and employment at some time between 20 and 24 weeks. Bed rest improves uterine blood flow and may increase birth weight up to 20%. Intercourse is generally discouraged when bed rest is recommended.

Monitoring a Multiple Pregnancy

Since preterm birth and growth disturbances are the major contributors to newborn death and disability in multiples, frequent obstetric visits and close monitoring of the pregnancy is needed.

Prenatal diagnosis by *chorionic villus sampling* can be done near the end of the first trimester to screen for Down syndrome and other genetic abnormalities. Amniocentesis is performed between 16 to 20 weeks. These procedures are complicated and difficult to perform in twins and triplets, and may not be possible in high order multiple pregnancies.

Many physicians perform cervical examinations every week or two beginning early in pregnancy to determine if the *cervix* is thinning or opening prematurely. If an exam or ultrasound shows that the cervix is thinning or beginning to dilate prematurely, a *cerclage*, or *suture* placed in the cervix, may prevent or delay premature dilatation.

Tocolytic agents are medications that may slow or stop premature labor. These medications are given in hospital “emergency” settings in an attempt to stop premature labor. It is important to attempt to delay delivery to minimize the risks of premature delivery.

Ultrasound examinations in the second trimester can identify some birth defects. Assessment of fetal growth by ultrasound every three to four weeks during the second half of pregnancy is commonly performed.

Every multiple pregnancy should be considered “high risk,” and obstetricians experienced with the management of multiple gestations should provide care. A neonatal intensive care unit nursery should be available to provide immediate and comprehensive support to premature newborns.

Cesarean Section

Vaginal delivery of twins may be safe in some circumstances. Many twins can be delivered vaginally if the presenting infant is in the head first position. Most triplets will be delivered by Cesarean section. Appropriate anesthesia and neonatal support are essential, whether delivery is performed vaginally or requires Cesarean section. Delivery of multiples requires planning by the entire medical team and availability of full intensive care support following birth.

Psychosocial Effects of Multiples on a Family

Although many women with a multiple pregnancy do well, their families may experience significant stress. If prolonged hospitalization is needed, arrangements must be made for work, home, and family care.

Even if medical problems are overcome and the infants survive without disability, the effect of multiple births on family life is profound and life-altering. The impact of a multiple birth clearly affects the parents, but also the babies, other siblings, and the extended family. Financial stresses may be overwhelming. Obvious additional costs include feeding, clothing, housing, and caring for multiple children.

Psychological counseling and support groups may provide a lifeline for the parents of multiples, who may feel isolated or depressed. Most physicians can provide appropriate referrals to a mental health professional or a support group. For more information, see the ASRM Patient Fact Sheet titled *Challenges of Parenting Multiples*.

CONCLUSION

The objective of infertility treatment is the birth of a healthy child. In a small percentage of patients, treatment results in multiple pregnancies that may place the mother and the babies at increased risk for an unhealthy outcome. Since multiple pregnancies and their complications are an inevitable risk of fertility therapies, education about these risks is crucial prior to treatment. Ultimately, prevention is the key to reducing the risk of multiple pregnancies.

NOTES

Let Us Know What You Think

Email your comments on this booklet to asmr@asmr.org. In the subject line, type "Attention: Patient Education Committee."

GLOSSARY

American Society for Reproductive Medicine (ASRM). A professional medical organization of approximately 9,000 health care specialists interested in reproductive medicine.

Amniocentesis. A procedure in which a small amount of amniotic fluid is removed through a needle from the fetal sac at about 16 weeks into a pregnancy. The fluid is studied for chromosomal abnormalities which may affect fetal development.

Amnion. Thin membrane that expands to enclose a developing fetus. This membrane (sac) holds the amniotic fluid that protects the developing fetus.

Assisted Reproductive Technology (ART). All treatments which include laboratory handling of eggs, sperm, and/or embryos. Some examples of ART are in vitro fertilization (IVF), gamete intrafallopian transfer (GIFT), pronuclear stage tubal transfer (PROST), tubal embryo transfer (TET), and zygote intrafallopian transfer (ZIFT).

Cerclage. Placement of a non-absorbable suture around an incompetent (weak) cervical opening in an attempt to keep it closed and thus prevent miscarriage. Also known as a cervical stitch.

Cerebral palsy. A disorder causing damage to one or more specific areas of the brain usually occurring during fetal development; before, during or shortly after birth; or infancy. Cerebral palsy is characterized by an inability to fully control motor function, particularly muscle control and coordination. Other problems that may arise are difficulties in feeding, bladder and bowel control, problems with breathing, skin disorders and learning disabilities.

Cervix. The lower narrow end of the uterus that connects the uterine cavity to the vagina.

Chorionic villus sampling (CVS). A procedure in which a small sample of cells is taken from the placenta early in a pregnancy for chromosomal testing.

Clomiphene citrate. An oral anti-estrogen drug used to induce ovulation in the female.

Diabetes. A condition due to abnormal production of insulin resulting in abnormally elevated blood glucose (sugar) levels.

Dizygotic. Di - two; zygote - fertilized egg. Two separate eggs fertilized by separate sperm in a single pregnancy. Fraternal twins.

Egg. The female sex cell produced by the ovaries, which, when fertilized by a male's sperm, produce embryos.

Embryo. The earliest stage of human development arising after the union of the sperm and egg (fertilization).

Fetus. An unborn child.

Follicle (ovarian). A fluid-filled sac located just beneath the surface of the ovary, containing an egg (oocyte) and cells that produce hormones. The sac increases in size and volume during the first half of the menstrual cycle and at

ovulation, the follicle matures and ruptures, releasing the egg. As the follicle matures, it can be visualized by ultrasound.

Genetic. Referring to inherited conditions, usually due to the genes located on the chromosomes.

Gestation. Pregnancy.

Gestational diabetes. Elevated blood sugar levels in the mother while she is pregnant. During pregnancy, the placenta normally produces hormones that antagonize insulin. With a multiple pregnancy, more of these hormones are produced and lead to a rise in the mother's blood sugar.

Gestational sac. The fluid-filled sac surrounding an embryo that develops within the uterine cavity. Ultrasound can detect the sac in the uterus at a very early stage of pregnancy.

Gonadotropins. Follicle stimulating hormone (FSH) and luteinizing hormone (LH). FSH and LH may be purified or synthetically produced to be used as ovulatory drugs.

Hormone. Substances secreted from organs of the body, such as the pituitary gland, adrenal gland, or ovaries, which are carried by a bodily fluid such as blood to other organs or tissues where the substances exert a specific action.

Infertility. Infertility is the result of a disease of the male or female reproductive tract which prevents the conception of a child or the ability to carry a pregnancy to delivery. The duration of unprotected intercourse with failure to conceive should be about 12 months or more before an investigation is undertaken, unless medical history, age, and physical findings dictate earlier evaluation and treatment.

In vitro fertilization (IVF). A method of assisted reproduction that involves combining an egg with sperm in a laboratory dish. If the egg fertilizes and begins cell division, the resulting embryo is transferred into the woman's uterus where it can implant in the uterine lining and further develop. IVF is generally performed in conjunction with medications that stimulate the ovaries to produce multiple eggs in order to increase the chances of successful fertilization and implantation. IVF bypasses the fallopian tubes and is often the treatment choice for women who have badly damaged or absent tubes.

Miscarriage. The naturally occurring expulsion of a nonviable fetus and placenta from the uterus; also known as spontaneous abortion or pregnancy loss.

Monozygotic. Mono - one; zygotic - fertilized egg. One egg fertilized by a single sperm that divides into two embryos. Maternal twins. Identical twins.

Multifetal pregnancy reduction. Also known as selective reduction. A procedure to reduce the number of fetuses in the uterus. As the risk of miscarriage (spontaneous abortion) and other problems increases with the number of fetuses present, this procedure may be performed in an attempt to prevent the entire pregnancy from aborting or delivering very prematurely.

Ovulation induction. The administration of hormone medications (ovulation drugs) that stimulate the ovaries to develop a follicle and ovulate.

Placenta. A disk-shaped vascular organ attached to the wall of the uterus and to the fetus by the umbilical cord. It provides nourishment to the fetus.

Preeclampsia. A disorder occurring during pregnancy that affects both the mother and the fetus. Preeclampsia is characterized by high blood pressure, swelling and protein found in the urine. This disorder, also known as toxemia, can restrict the flow of blood to the placenta.

Respiratory distress syndrome (RDS). A lung disease that affects premature infants and causes increasing difficulty in breathing.

Singleton. Offspring (child) born singly.

Sperm. The male reproductive cells that fertilize a woman's egg. The sperm head carries genetic material (chromosomes), the midpiece produces energy for movement, and the long, thin tail wiggles to propel the sperm.

Superovulation. The administration of fertility medications in a manner intended to achieve development and ovulation of multiple ovarian follicles. Superovulation is often combined with intrauterine insemination as an infertility treatment.

Suture. Thread used to close an incision made during surgery. It is generally absorbable or self-dissolving.

Tocolytic agents. Medications that may slow or stop premature labor.

Toxemia. See Preeclampsia.

Ultrasound. Also called sonogram. A picture of internal organs produced by high frequency sound waves viewed as an image on a video screen; used to monitor growth of ovarian follicles, to retrieve eggs, and to monitor a fetus or pregnancy. Ultrasound examinations can be performed either abdominally or vaginally.

Uterus (Womb). The hollow, muscular organ in the pelvis where an embryo implants and grows during pregnancy. The lining of the uterus, called the endometrium, produces the monthly menstrual blood flow when there is no pregnancy.

For a list of additional reading materials, contact the ASRM administrative office at 1209 Montgomery Highway, Birmingham, Alabama 35216-2809; (205) 978-5000.

Booklets available for purchase through the
American Society for Reproductive Medicine Patient Information Series:

- ___ *Abnormal Uterine Bleeding (1996)*
- ___ *Adoption (1996)*
- ___ *Age and Fertility (2003)*
- ___ *Assisted Reproductive Technologies (2003)*
- ___ *Birth Defects of the Female Reproductive System (1993)*
- ___ *Donor Insemination (1995)*
- ___ *Ectopic Pregnancy (1996)*
- ___ *Endometriosis (1995)*
- ___ *Endometriosis (Spanish Translation) (2003)*
- ___ *Fertility After Cancer Treatment (1995)*
- ___ *Hirsutism and Polycystic Ovarian Syndrome (2003)*
- ___ *Husband Insemination (1995)*
- ___ *Infertility: An Overview (2003)*
- ___ *Infertility: An Overview (Spanish Translation) (1996)*
- ___ *Infertility: Coping and Decision Making (1995)*
- ___ *Laparoscopy and Hysteroscopy (1995)*
- ___ *Male Infertility and Vasectomy Reversal (1995)*
- ___ *Miscarriage (1995)*
- ___ *Multiple Pregnancy and Birth: Twins, Triplets, & Higher Order Multiples (2004)*
- ___ *Ovulation Detection (1995)*
- ___ *Ovulation Drugs (2000)*
- ___ *Pelvic Pain (1997)*
- ___ *Pregnancy After Infertility (1997)*
- ___ *Premenstrual Syndrome (PMS) (1997)*
- ___ *Third Party Reproduction (Donor Eggs, Donor Sperm, Donor Embryos,
& Surrogacy) (1996)*
- ___ *Tubal Factor Infertility (1995)*
- ___ *Unexplained Infertility (1998)*
- ___ *Uterine Fibroids (1997)*

For copies, ask your physician or contact the ASRM at the address below.

AMERICAN SOCIETY FOR REPRODUCTIVE MEDICINE

1209 Montgomery Highway • Birmingham, Alabama 35216-2809
(205) 978-5000 • asrm@asrm.org • www.asrm.org



AMERICAN SOCIETY FOR REPRODUCTIVE MEDICINE

1209 Montgomery Highway
Birmingham, Alabama 35216-2809
(205) 978-5000 • asrm@asrm.org • www.asrm.org